

THE RING OF ARITHMETICAL FUNCTIONS WITH UNITARY CONVOLUTION: THE $[n]$ -TRUNCATION

JAN SNELLMAN

ABSTRACT. We study a certain truncation $\mathcal{A}_{[n]}$ of the ring of arithmetical functions with unitary convolution, consisting of functions vanishing on arguments $> n$. The truncations $\mathcal{A}_{[n]}$ are artinian monomial quotients of a polynomial ring in finitely many indeterminates, and are isomorphic to the “artinified” Stanley-Reisner ring $\mathbb{C}[\overline{\Delta([n])}]$ of a simplicial complex $\Delta([n])$.

1. INTRODUCTION

The set $\mathcal{A}$ of all complex sequences $(c_i)_{i=1}^\infty$ is in a natural way a complex vector space. There are several ways an algebra structure can be introduced on this vector space: the most intuitive way are through the so-called *regular convolutions* of Narkiewicz [14]. Among those, the *Dirichlet convolution* and the *unitary convolution* are the best known specimen. They are in a way the two extremes, in that they form the maximal and minimal elements in a certain natural partial order on the set of all regular convolutions.

The ring given by Dirichlet convolution is isomorphic to the formal power series ring on countably many indeterminates, hence a domain; in fact, it is a UFD, as proved by Cashwell and Everett [4]. As a contrast, the ring given by unitary convolution is an epimorphic image of the formal power series ring on countably many indeterminates: it has zero-divisors and nilpotent element. In [17] we gave a conjectural characterization of the zero-divisors. We also established some divisorial properties: a given element can have factorizations (into irreducibles) of different length, but there is always a bound for those lengths.

Let us denote by $\mathcal{A}_{[n]}$ the subset of $\mathcal{A}$ consisting of those sequences $(c_i)_{i=1}^\infty$ for which $c_i = 0$ when $i > n$. Restricting any convolution product on $\mathcal{A}$, and modifying it so that the product of two elements in $\mathcal{A}_{[n]}$ stays in $\mathcal{A}_{[n]}$, we get a new algebra, which is an artinian monomial quotient of the original one. These quotients, for the case of the Dirichlet convolution, were studied in [16]. It turned out that in this case

Date: August 23, 2002.

1991 Mathematics Subject Classification. 13F55, 11A25.

Key words and phrases. Arithmetical functions, unitary convolution, simplicial complexes, Stanley-Reisner rings.

the defining ideals are *strongly stable*, hence we can apply the Eliahou-Kervaire resolution [6] to get information about various homological invariants.

In this article, we study the algebra $\mathcal{A}_{[n]}$ equipped with the multiplication given by the unitary convolution. It turns out that the defining ideals, apart from being monomial ideals, are *almost square-free*, i.e. they are the sum of a square-free ideal and the ideal generated by the squares of the indeterminates. The ring is therefore the quotient of $\frac{\mathbb{C}[x_1, \dots, x_r]}{(x_1^2, \dots, x_r^2)}$ by a square-free monomial ideal, hence it is the *artinified Stanley-Reisner ring* $\mathbb{C}[\overline{\Delta}]$, in the sense of Sköldbberg [15], of a simplicial complex Δ . In short, the artinified Stanley-Reisner ring is like the indicator algebra $\mathbb{C}\{\Delta\}$ (see for instance [1] for a definition) but it is a quotient of $\frac{\mathbb{C}[x_1, \dots, x_r]}{(x_1^2, \dots, x_r^2)}$ rather than a quotient of the exterior algebra, so it is commutative rather than skew-commutative.

The simplicial complex Δ is easy to describe: the underlying vertex set consists of all prime powers $p^a \leq n$, and a set $\{q_1, \dots, q_s\}$ of such prime powers is a simplex in Δ if and only if the q_i 's are relatively prime, and their product is $\leq n$. A figure is supposedly worth a thousand words, so the reader may want to look at Figure 3 on page 8, where the complex for $n = 10$ is displayed. Understanding this complex, particularly for large n , is the purpose of this article. In particular, we determine the dimension and the largest non-vanishing homology of Δ , and show that all homology groups are torsion-free, by exhibiting a (non-pure) shelling. We also give an asymptotic estimate of the socle dimension of $\mathcal{A}_{[n]}$.

In [18] a study was made of the truncation of $\mathcal{A}$ to some subset $V \subset \mathbb{N}^+$ which is closed under taking unitary divisors. The results obtained for the corresponding algebras $\mathcal{A}_V$, in particular for finite V , will be of use for us in our study of the special case $V = [n] = \{1, \dots, n\}$, so this article starts with a review of the pertinent definitions and results from [18].

1.1. Acknowledgement. I used the GAP-package *Simplicial Homology* [5] by Frank Heckenbach, Jean-Guillaume Dumas, Dave Saunders, and Volkmar Welker to calculate the homology of Δ . Having noted that the homology was torsion-free, I wrote a small GAP [8] programme to check if it was lex-shellable: it was, and it was then easy to prove that. I also benefitted from the programme Macaulay 2 [10] to calculate Poincaré-Betti series of $\mathcal{A}_{[n]}$.

2. PRELIMINARIES

This article is a direct continuation of [18], from which we recall some definitions and basic results.

2.1. The ring of arithmetical functions with unitary convolution. Let $\mathbb{N}$ denote the non-negative integers and $\mathbb{N}^+$ the positive integers. Let $\mathbb{P}$ denote the set of primes, with $p_i \in \mathbb{P}$ the i 'th prime. Let $\mathbb{PP}$ denote the set of prime powers. $\mathcal{A}$ denotes the set of *arithmetical functions*, i.e. functions $\mathbb{N}^+ \rightarrow \mathbb{C}$. It is a $\mathbb{C}$ -vector space under point-wise addition and multiplication by scalars, and it has a natural topology given by the norm

$$|f| = \frac{1}{\min \text{supp}(f)},$$

where

$$\text{supp}(f) = \{k \in \mathbb{N}^+ \mid f(k) \neq 0\}.$$

$\mathcal{A}$ becomes an associative, commutative $\mathbb{C}$ -algebra under *unitary convolution*

$$(f \oplus g)(n) = \sum_{d \parallel n} f(d)g(n/d) = \sum_{d \oplus m = n} f(d)g(m), \quad (1)$$

where the unitary multiplication for positive integers is defined by

$$d \oplus m = \begin{cases} dm & \text{if } \gcd(d, m) = 1 \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

and where we write $d \parallel n$ (or sometimes $d \leq_{\oplus} n$) when d is a unitary divisor of n , i.e. when $n = d \oplus m$ for some m . For any $k \in \mathbb{N}^+$, e_k denotes the characteristic function on $\{k\}$. Then e_1 is the multiplicative identity, the set of all e_k is a Schauder basis for $\mathcal{A}$, and

$$e_a \oplus e_b = e_{a \oplus b} = \begin{cases} e_{ab} & \text{if } \gcd(a, b) = 1 \\ 0 & \text{otherwise} \end{cases}$$

so $\{e_k \mid k \in \mathbb{PP}\}$ generates a dense subalgebra of $\mathcal{A}$.

Let $Y = \{y_{i,j} \mid i, j \in \mathbb{N}^+\}$ be a doubly infinite set of indeterminates, and let $[Y]$ be the free abelian monoid on Y . Let $\mathcal{M} \subset [Y]$ consist of those monomials in the $y_{i,j}$'s that are *separated*, i.e. can be written $y_{i_1, j_1} \cdots y_{i_r, j_r}$ with $i_1 < i_2 < \cdots < i_r$. Then $\mathcal{M}$ can be regarded as a monoid-with-zero, with the multiplication

$$m_1 \cdot m_2 = \begin{cases} m_1 m_2 & \text{if } m_1 m_2 \in \mathcal{M} \\ 0 & \text{if } m_1 m_2 \notin \mathcal{M} \end{cases}$$

and

$$\begin{aligned} \Phi : \mathcal{M} &\rightarrow \mathbb{N}^+ \\ y_{i_1, j_1} \cdots y_{i_r, j_r} &\mapsto p_{i_1}^{j_1} \cdots p_{i_r}^{j_r} \end{aligned} \quad (3)$$

is a bijection which is a monoid-with-zero isomorphism, if $\mathbb{N}^+$ is regarded as a monoid-with-zero with unitary multiplication. From this

follows that

$$\mathcal{A} \simeq \mathbb{C}[[\mathcal{M}]] \simeq \frac{\mathbb{C}[[Y]]}{J} \quad (4)$$

where $\mathbb{C}[[\mathcal{M}]]$ and $\mathbb{C}[[Y]]$ are the generalized power series rings on $\mathcal{M}$ and $[Y]$, respectively (so $\mathbb{C}[[Y]]$ is the power series ring on bi-infinitely many variables) and J is the smallest closed ideal of $\mathbb{C}[[Y]]$ which contains all $y_{i,j}y_{i,k}$.

2.2. General truncations. For any $V \subseteq \mathbb{N}^+$, $\mathcal{A}_V \subseteq \mathcal{A}$ is the $\mathbb{C}$ -sub vector space of functions supported on V . With the modified multiplication

$$(f \oplus g)(n) = \sum_{d \oplus_V m = n} f(d)g(m) \quad (5)$$

$$d \oplus_V m = \begin{cases} dm & \text{if } \gcd(d, m) = 1 \text{ and if } dm \in V \\ 0 & \text{otherwise} \end{cases}$$

it becomes a $\mathbb{C}$ -algebra, but in general not a sub-algebra of $\mathcal{A}$; it is a sub-algebra if and only if

$$a, b \in V \implies a \oplus b \in V \cup \{0\}.$$

If V contains all unitary divisors of its elements, then the restriction map $\mathcal{A} \rightarrow \mathcal{A}_V$, which is always a vector space epimorphism, is an algebra epimorphism. In particular, if n is a positive integer, then the set $[n] = \{1, 2, \dots, n\} \subset \mathbb{N}^+$ has this property. If we denote the kernel of the restriction map by $\mathfrak{S}_V$, then

Theorem 2.1. *The set*

$$M_V = \{e_k | k \notin V, \text{ but } d \in V \text{ for all proper unitary divisors } d \text{ of } k\} \quad (6)$$

form a minimal generating set of an ideal I_V whose closure is $\mathfrak{S}_V$.

2.3. Finite truncations. Now suppose that V has this property, and is finite. Put

$$Y(V) = \{y_{i,j} | p_i^j \in V \cap \mathbb{P}\mathbb{P}\} \quad (7)$$

From [18] we have that

$$\mathcal{A}_V \simeq \frac{\mathbb{C}[Y(V)]}{A_V + B_V + C_V} \quad (8)$$

where

$$\begin{aligned} A_V &= \langle y_{i,j}^2 | y_{i,j} \in Y(V) \rangle \\ B_V &= \langle y_{i,j}y_{i,k} | y_{i,j}, y_{i,k} \in Y(V) \rangle \\ C_V &= \langle y_{i_1,j_i} \cdots y_{i_r,j_r} | y_{i_\ell,j_\ell} \in Y(V), i_1 < i_2 < \cdots < i_r, p_{i_1}^{j_i} \cdots p_{i_r}^{j_r} \notin V \rangle \end{aligned} \quad (9)$$

We also have that

$$\mathcal{A}_V \simeq \mathbb{C}[\overline{\Delta(V)}] \quad (10)$$

where $\Delta(V)$ is the simplicial complex on the vertex set $V \cap \mathbb{P}\mathbb{P}$ given by

$$\sigma = \{p_{i_1}^{j_1}, \dots, p_{i_r}^{j_r}\} \in \Delta(V) \iff p_{i_1}^{j_1} \cdots p_{i_r}^{j_r} \in V \quad (11)$$

and where $\mathbb{C}[\overline{\Delta(V)}]$ is the “Artinified Stanley-Reisner ring” [15] on $\Delta(V)$: it is the artinian commutative $\mathbb{C}$ -algebra with a $\mathbb{C}$ -basis

$$\{e_\sigma \mid \sigma \in \Delta(V)\},$$

and multiplication

$$e_\sigma e_\tau = \begin{cases} e_{\sigma \cup \tau} & \text{if } \sigma \cap \tau = \emptyset \\ 0 & \text{if } \sigma \cap \tau \neq \emptyset \end{cases} \quad (12)$$

Thus, as a graded $\mathbb{C}$ -vector space $\mathbb{C}[\overline{\Delta(V)}]$ is isomorphic to $\mathbb{C}\{\Delta(V)\}$, the indicator algebra on $\Delta(V)$ [1], and as a cyclic $\mathbb{C}[Y(V)]$ -module it is isomorphic to the quotient of the ordinary Stanley-Reisner ring $\mathbb{C}[\Delta(V)]$ by the ideal generated by the squares of all variables.

2.4. The $[n]$ -truncation. This article is devoted to the case $V = [n] = \{1, 2, \dots, n\}$, for which the ring $\mathcal{A}_V$ and the simplicial complex $\Delta(V)$ have some interesting properties.

3. SOME SPECIAL ARITHMETICAL FUNCTIONS

We’ll make use of the following special arithmetical functions: For a positive integer n , $\pi(n)$ is the number of primes $\leq n$, and $\pi'(n)$ the number of prime powers $\leq n$. Let $\omega(n)$ denotes the number of distinct prime factors of n , and let (for $k \leq 0$) $\pi_k(n)$ be the number of positive integers $\leq n$ with k distinct prime factors, i.e.

$$\sum_{k=0}^{\infty} \pi_k(n) t^k = \sum_{j=1}^n t^{\omega(j)}.$$

By $\ell(n)$ we mean the unique integer such that

$$\prod_{i=1}^{\ell(n)} p_i \leq n < \prod_{i=1}^{\ell(n)+1} p_i, \quad (13)$$

We define $v(1) = v(2) = -1$, and for $n \geq 3$, $v(n)$ as the unique integer such that

$$\prod_{j=2}^{v(n)+1} p_j \leq n < \prod_{j=2}^{v(n)+2} p_j \quad (14)$$

It follows that $v(n) = \ell(2n) - 2$. The values of $\ell(n)$ and $v(n)$ for small n are tabulated below.

n	1	2	3	4	5	6	...	15	...	30	...	105	...	210
$\ell(n)$	0	1	1	1	1	2	...	2	...	3	...	3	...	4
$v(n)$	-1	-1	0	0	0	0	...	1	...	1	...	2	...	2

4. THE STRUCTURE OF $Y([n])$

We now determine the structure of $Y(V)$ for $V = [n]$.

Definition 4.1. For all positive integers n, i , let

$$\begin{aligned}\lambda_i^{[n]} &= \max \{ j \mid p_i^j \leq n \} \\ \boldsymbol{\lambda}^{[n]} &= (\lambda_1^{[n]}, \lambda_2^{[n]}, \dots)\end{aligned}\tag{15}$$

We may regard $\boldsymbol{\lambda}^{[n]}$ as a partition, since $\lambda_1^{[n]} \geq \lambda_2^{[n]} \geq \dots$. We have that

$$Y([n]) = \left\{ y_{i,j} \in Y \mid j \leq \lambda_i^{[n]} \right\}\tag{16}$$

Note that $\max(\{ i \mid y_{i,1} > 0 \}) = \pi'(n)$.

Example 4.2. If $n = 30$, then

$$\boldsymbol{\lambda}^{[30]} = (4, 3, 2, 1, 1, 1, 1, 1, 1, 1),$$

and the variables $Y(V_{[30]})$ can be visualised as in figure 1.

	2	3	5	7	11	13	17	19	23	29
$y_{1,4}$										
$y_{1,3}$	$y_{2,3}$									
$y_{1,2}$	$y_{2,2}$	$y_{3,2}$								
$y_{1,1}$	$y_{2,1}$	$y_{3,1}$	$y_{4,1}$	$y_{5,1}$	$y_{6,1}$	$y_{7,1}$	$y_{8,1}$	$y_{9,1}$	$y_{10,1}$	

FIGURE 1. $Y(\boldsymbol{\lambda}^{[30]})$

Remark 4.3. For a fixed i , as $n \rightarrow \infty$ we have that asymptotically $\lambda_i^{[n]} \sim \frac{\log(n)}{\log(p_i)}$. Thus, for both n and i large we get that $\lambda_i^{[n]} \sim \frac{\log(n)}{\log(i \log(i))}$. This relation is illustrated in the graph figure 2, which shows (the start of) $\boldsymbol{\lambda}^{(10^{50})}$.

5. THE PRESENTATION OF $\mathcal{A}_{[n]}$

Having determined the indeterminates occuring in $Y([n])$, we'll consider the defining ideal $A_{[n]} + B_{[n]} + C_{[n]}$ of $\mathcal{A}_{[n]} \simeq \frac{\mathbb{C}[Y([n])]}{(A_{[n]} + B_{[n]} + C_{[n]})}$. Recall that (3) defines a bijection Φ between $\mathcal{M}$ and $\mathbb{N}^+$.

Theorem 5.1. *Let $V = [n]$. Then*

1. The minimal generators of $C_{[n]}$ correspond under Φ with those e_k for which
 - $k > n$,
 - all proper unitary divisors of k are $\leq n$,
 - $k = \prod_{i=1}^r p_i^{a_i}$ with $r \leq \pi'(n)$ and $0 \leq a_i \leq \lambda_i^{[n]}$ for $1 \leq i \leq r$.
2. $A_{[n]} + B_{[n]} + C_{[n]}$ is a strongly $N + 1$ -multi-stable monomial ideal in $\mathbb{C}[Y([n])]$. By this, we mean the following: first, let N denote the number of components > 1 in $\lambda^{[n]}$. Group the variables $y_{i,j}$ in column i into one group, for $1 \leq i \leq N$, and the remaining variables into one last group. Order the variables in each group using Φ . Then for any monomial $m \in A_{[n]} + B_{[n]} + C_{[n]}$, if y, y' belong to the same group, $\Phi(y) < \Phi(y')$, and $y \mid m$, then $m' = \frac{y'}{y}m \in A_{[n]} + B_{[n]} + C_{[n]}$.


Proof. 1. Follows from Theorem 2.1 and (8) and (9).

2. If $m \in A_{[n]} + B_{[n]}$ then it contains two variables “from the same column”. Since y, y' belong to the same group, m' must also contain two variables “from the same column”; thus $m' \in A_{[n]} + B_{[n]}$.

If m is separated, then so is m' , and $\Phi(m) < \Phi(m')$ so if in addition $m \in C_{[n]}$ then $m' \in C_{[n]}$.

□

FIGURE 2. $\lambda^{(10^{50})}$



The following lemma gives a reasonably efficient way of calculating $C_{[n]}$.

Lemma 5.2. *The minimal generators of $C_{[n]}$ are contained in the set*

$$\mathcal{M} \cap \{ y_{i,j} \Phi^{-1}(k) | p_i^j \leq n, 1 \leq k \leq n \} \setminus \Phi^{-1}(\{1, 2, \dots, n\}) \quad (17)$$

Proof. Take $m = m_0 \in \mathcal{M} \cap Y([n])$, $\Phi(m) > n$. Thus m is a separated monomial in variables $y_{i,j}$ with $\Phi(y_{i,j}) \leq n$. Hence there is some y_{i_1,j_1} with $\Phi(y_{i_1,j_1}) \leq n$ which divides m , say $m = y_{i_1,j_1} m_1$, $m_1 \in \mathcal{M} \cap Y([n])$. If $\Phi(m_1) \leq n$ we are done, otherwise m_1 , being a separated monomial in variables $y_{i,j}$ with $\Phi(y_{i,j}) \leq n$, can be written $m_1 = y_{i_1,j_2} m_2$, et cetera. $\square$

Example 5.3. For $n = 10$, $\lambda^{(10)} = (3, 2, 1, 1)$, and

$$Y(V_{[10]}) = \{y_{1,1}, y_{1,2}, y_{1,3}, y_{2,1}, y_{2,2}, y_{3,1}, y_{4,1}\}$$

$$A_{10} = (y_{1,1}y_{1,1}, y_{1,2}y_{1,2}, y_{1,3}y_{1,3}, y_{2,1}y_{2,1}, y_{2,2}y_{2,2}, y_{3,1}y_{3,1}, y_{4,1}y_{4,1})$$

$$B_{10} = (y_{1,1}^2, y_{1,1}^2, y_{1,2}^2, y_{2,1}^2)$$

$$C_{10} = (y_{1,1}y_{2,2}, y_{1,1}y_{4,1}, y_{2,2}y_{3,1}, y_{1,2}y_{2,1}, y_{1,3}y_{2,1}, y_{2,1}y_{3,1}, y_{2,1}y_{4,1}, \\ y_{1,2}y_{2,2}, y_{1,2}y_{3,1}, y_{1,2}y_{4,1}, y_{1,3}y_{3,1}, y_{1,3}y_{4,1}, y_{2,2}y_{4,1}, y_{1,3}y_{2,2}, y_{3,1}y_{4,1}, \\ y_{1,1}y_{2,1}y_{3,1}, y_{1,1}y_{2,1}y_{4,1}, y_{1,1}y_{2,2}y_{3,1}, y_{1,1}y_{3,1}y_{4,1})$$


where the last four generators are superfluous.

In Corollary 8.3, we show that the maximal degree of a minimal generator of $C_{[n]}$ is $v(n)$.

6. PROPERTIES OF $\Delta([n])$

As an example of $\Delta([n])$, $\Delta([10])$ is shown in Figure 3.

FIGURE 3. $\Delta([10])$



6.1. Simple properties. Let $r = \pi'([n])$. Clearly $\Delta([n])$ has r vertices.

Lemma 6.1. *$\Delta([n])$ consists of some isolated vertices, and of one large component containing everything else.*

Proof. If the vertex p^a is not isolated, but connected to q^b , we can assume that $p > q$. Furthermore, since $\{p^a, q^b\} \in \Delta([n])$, it follows that $p^a q^b \leq n$, so $p^a q \leq n$, hence p^a is connected to q . If $q > 2$ then $p > q > 2$ so $\{p^a, 2\} \in \Delta([n])$. Thus p^a is connected to 2. $\square$

Lemma 6.2. $\dim \Delta([n]) = \ell(n) - 1$.

Proof. The maximal cardinality s of $(q_1, \dots, q_s) \in \Delta([n])$ is the maximal s such that

$$\prod_{i=1}^s q_i \leq n,$$

with $q_i \in \mathbb{P}$ pair-wise relatively prime. Clearly, the best we can do is to take the first s prime numbers, so $s = \ell(n)$. Hence $\dim \Delta([n]) = s - 1 = \ell(n) - 1$. $\square$

Lemma 6.3. *The f -vector of $\Delta([n])$ is*

$$(f_{-1}, f_0, f_1, \dots, f_{\ell(n)-1}) = (\pi_0(n), \pi_1(n), \pi_2(n), \dots, \pi_{\ell(n)}(n)) \quad (18)$$

Proof. We have that f_i is the number of simplicies $\sigma \in \Delta([n])$ of dimension i , i.e. of cardinality $i + 1$. Such a $\sigma = \{p_{j_1}^{a_1}, \dots, p_{j_{i+1}}^{a_{i+1}}\}$ correspond to $k = p_{j_1}^{a_1} \cdots p_{j_{i+1}}^{a_{i+1}}$ with $\omega(k) = i + 1$. There are $\pi_{i+1}(n)$ such simplices, so $f_i = \pi_{i+1}(n)$. $\square$

Lemma 6.4. *Let $v(n)$ be defined by (14). Then*

- (i) *The homological degree¹ of $\Delta([n])$ is $v(n)$.*
- (ii) *The maximal homological degree of all $\Delta([n])_U$, as U ranges among the non-empty subsets of $[n]$, is $v(n)$.*

Sketch of proof. When $n = N = \prod_{j=2}^s p_j$, we have that

$$\{p_1, \dots, p_s\} \notin \Delta([n]), \quad \text{but } \forall i : \{p_1, \dots, \widehat{p_i}, \dots, p_s\} \in \Delta([n])$$

so we get $(s - 1)$ -homology. As n increases and reaches

$$2N = \prod_{j=1}^s p_j,$$

this homology is killed off, but already when

$$n = N \frac{p_{s+1}}{p_s} = p_{s+1} \prod_{j=2}^{s-1} p_j$$

new $s - 1$ homology appears, since all $(s - 1)$ -subsets of $\{p_1, \dots, p_{s-1}, p_{s+1}\}$ belong to $\Delta([n])$, whereas the whole set doesn't. Filling in this homology, i.e. increasing n to $p_1 \cdots p_{s-1} \cdot p_{s+1}$, we have introduced new homology already at

$$n = p_2 \cdots p_{s-1} \cdot p_{s+2}$$

¹The maximal i such that the i 'th reduced homology group (with coefficients in $\mathbb{Z}$) of $\Delta([n])$ is non-zero.

(use the fact that there is a prime number between q and $2q$ for all q), and so on, until we reach

$$n = \prod_{j=2}^{s+1} p_j,$$

where s -homology occurs. $\square$

The asymptotic growth of $\ell(n)$ as $n \rightarrow \infty$ is very slow, as the following lemma shows. Thus $\dim \Delta([n]) = \ell(n)$ and $v(n) = \ell(2n) - 2$, the homological degree, grows very slowly with n .

Lemma 6.5. *There are positive real constants A, B, C, D such that for all n ,*

$$A \frac{\log(n)}{W(A \log(n))} < \ell(n) < B \frac{\log(n)}{W(B \log(n))} \quad (19)$$

where $W(z)$ denotes the real-valued principal branch of the Lambert W -function, defined as the root of $W(z) \exp(W(z)) = z$.

Proof. There are positive real constants² A_1, A_2 such that

$$A_1 x < \sum_{p \leq x} \log(p) < A_2 x,$$

hence

$$A_1 p_n < \sum_{i=1}^n \log(p_i) < \sum_{i=1}^{n+1} \log(p_i) < A_2 p_{n+1}.$$

There are constants³ B_1, B_2 such that

$$B_1 n \log(n) < p_n < p_{n+1} < B_2 (n+1) \log(n+1).$$

Put $m = \ell(n)$. Then

$$\sum_{i=1}^m \log(p_i) \leq \log(n) < \sum_{i=1}^{m+1} \log(p_i),$$

so

$$A_1 B_1 m \log(m) < \log(n) < A_2 B_2 (m+1) \log(m+1).$$

We claim that

$$C m \log(m) = \log(n) \quad (20)$$

has the solution

$$m = \frac{\log(n)}{C W(\frac{\log(n)}{C})}.$$

From this claim, the assertion follows by monotonicity.

²See [11]

³See [11]

Putting $\log(n) = z$, $\log(m) = a(z)$, (20) becomes

$$Ce^{a(z)}a(z) = z,$$

which has a solution $a(z) = W(z/C)$. Hence

$$m = \exp(a(z)) = \exp(W(z/C)) = \frac{z/C}{W(z/C)} = \frac{\log n}{CW(\frac{\log(n)}{C})}.$$

□

6.2. The h -vector. Using a result by Fröberg, we have that

$$\mathbb{C}[\Delta([n])](t) = \mathbb{C}[\overline{\Delta(n)}](\frac{t}{1-t}) = \frac{h_0 + h_1t + h_2t^2 + \cdots + h_{\ell(n)}t^{\ell(n)}}{(1-t)^{\ell(n)}} \quad (21)$$

where $h_0 + h_1 + \cdots + h_{\ell(n)} \neq 1$. The vector $(h_0, \dots, h_{\ell(n)})$ is called the h -vector of $\Delta([n])$. It can be expressed in terms of the f -vector as follows (see [19]).

Lemma 6.6. *The h -vector of $\Delta([n])$ is*

$$(h_0, h_1, \dots, h_{\ell(n)}) \quad (22)$$

where

$$\begin{aligned} h_k &= \sum_{i=0}^k (-1)^{k-i} \binom{\ell(n)-i}{k-i} f_{i-1} \\ &= \sum_{i=0}^k (-1)^{k-i} \binom{\ell(n)-i}{k-i} c_{n,i} \end{aligned} \quad (23)$$

with the convention that $f_{-1} = c_{n,0} = 1$.

In particular,

$$\begin{aligned} h_0 &= 1 \\ h_1 &= -\ell(n) + \pi'(n) \\ h_2 &= \binom{\ell(n)}{2} - (\ell(n) - 1)\pi'(n) + c_{n,2} \end{aligned} \quad (24)$$

Example 6.7. Let $\Delta = \Delta([10])$. Then Δ is 2-dimensional, looks like Figure 3, and has f -vector $(f_{-1}, f_0, f_2) = (1, 7, 2)$ and h -vector $(h_0, h_1, h_2) = (1, 5, -4)$. Furthermore

$$\mathbb{C}[\overline{\Delta}](t) = 1 + 7t + 2t^2,$$

and

$$\mathbb{C}[\Delta](t) = 1 + t \frac{t}{1-t} + 2 \frac{t^2}{(1-t)^2} = \frac{1 + 5t - 4t^2}{(1-t)^2}.$$

6.2.1. *The h_2 coefficient.* Clearly, $h_1 > 0$ for $n > 2$. Furthermore, we have (see [20])

$$\pi_k(x) \sim \frac{x}{\log x} \frac{(\log_2 x)^{k-1}}{(k-1)!} \quad x \rightarrow \infty \quad (25)$$

Thus we can write (somewhat sloppily)

$$\begin{aligned} h_2 &= \binom{\ell(n)}{2} - (\ell(n) - 1)\pi'(n) + c_{n,2} \\ &\approx \left(C \frac{\log(n)}{W(C \log(n))} \right)^2 - \left(C \frac{\log(n)}{W(C \log(n))} \right) \frac{n}{\log n} + \frac{n}{\log n} \log_2 n \\ &= -\frac{Cn}{W(C \log n)} + \frac{C^2(\log n)^2}{W(C \log n)^2} + \frac{n \log 2}{(\log n)^2} \end{aligned}$$

We claim that $h_2 < 0$ for large n , i.e. that

$$n > \frac{C(\log n)^2}{W(C \log n)} + n \frac{\log 2}{C} \frac{W(C \log n)}{(\log n)^2}$$

for large n . Since $n \gg C^2(\log n)^2$, we need only show that


$$\frac{W(C \log n)}{(\log n)^2} \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (26)$$

However, since $\ell(n) \rightarrow \infty$ as $n \rightarrow \infty$, (19) shows that

$$\frac{W(C \log n)}{C \log n} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

hence (26) follows.

6.3. Related questions. We display below how h_2 varies with n . Note that h_2 has local maxima when $\ell(n) < \ell(n+1)$, i.e. when n is of the form $n = -1 + \prod_{k=1}^j p_k$. Is the h_2 coefficient negative for all n ?



One can also ask: what is the sign (and magnitude) of the h_k coefficient, for very large n ?

6.4. Shellability. Recall that the maximal (with respect to inclusion) faces of a simplicial complex Δ are called *facets*, and that a simplicial complex is *pure* if all its facets have the same dimension (which is then also the dimension of the simplicial complex itself). For a face $\sigma \in \Delta$, we let $\bar{\sigma} = 2^\sigma$, the set of all subsets of σ , including σ itself, and the empty set.

Björner and Wachs [2, 3] defines Δ to be *shellable* if its facets can be arranged in linear order $F_1, F_2, \dots, F_t$ in such a way that the subcomplex $\cup_{i=1}^{k-1} \overline{F_i} \cap \overline{F_k}$ is pure and of dimension $\dim F_k - 1$ for all $k = 2, \dots, t$. They proved the following [2, Lemma 2.3]:

Lemma 6.8 (Björner-Wachs). *An order $F_1, F_2, \dots, F_t$ of the facets of Δ is a shelling if and only if for every i and k with $1 \leq i \leq k$ there is a j with $1 \leq j < k$ and an $x \in F_k$ such that $F_i \cap F_k \subseteq F_j \cap F_k = F_k \setminus \{x\}$.*

We define the *lexicographic order* on finite subsets of $\mathbb{P}\mathbb{P}$ in the following way: if

$$\sigma = a_1, \dots, a_r \subset \mathbb{P}\mathbb{P} \quad \text{with } 1 \leq a_1 < a_2 < \dots < a_r,$$

$$\tau = \{b_1, \dots, b_s\} \in \Delta([n]), \quad \text{with } 1 \leq b_1 < b_2 < \dots < b_s,$$

then $\sigma \geq_{lex} \tau$ if either $\tau = \emptyset$ or $a_1 > b_1$ or $\sigma \setminus \{a_1\} \geq_{lex} \tau \setminus \{b_1\}$. Note that $\geq_{lex}$ is a *Boolean term-order* in the sense of MacLagan [13], so

$$\sigma \geq_{lex} \tau \implies \sigma \cup \{w\} \geq_{lex} \tau \cup \{w\} \geq_{lex} \tau. \quad (27)$$

Theorem 6.9. $\Delta([n])$ is shellable; the lexicographic order on the facets is a shelling order.

Proof. Let $F_1, \dots, F_t$ be the facets in $\Delta([n])$ ordered lexicographically. Using the previous lemma, we'll show that this is a shelling. So pick $1 \leq i < k \leq t$, and let

$$\begin{aligned} F_i \cap F_k &= \{a_1, \dots, a_r\}, & 1 \leq a_1 < a_2 < \dots < a_r \leq n \\ F_i &= \{a_1, \dots, a_r\} \cup \{b_1, \dots, b_t\}, & 1 \leq b_1 < b_2 < \dots < b_t \leq n \\ F_k &= \{a_1, \dots, a_r\} \cup \{c_1, \dots, c_s\}, & 1 \leq c_1 < c_2 < \dots < c_s \leq n \end{aligned}$$

We always have that

$$r = |F_i \cap F_k| < |F_k| = r + s,$$

since by definition, no facet is contained in another facet. If

$$|F_k| - |F_i \cap F_k| = 1,$$

then we are done, by taking $j = i$.

So suppose that

$$|F_k| - |F_i \cap F_k| \geq 2.$$

This means that $s \geq 2$. Since $i < k$, $F_i >_{lex} F_k$. We distinguish two cases: some c_v is a prime power which is not a prime (case 1), or all c_v 's are prime (case 2).

Case 1: There is some c_v which is not a prime, thus $c_v = p^\delta$ with $\delta > 1$, p a prime. Put

$$G = (F_k \cup \{p\}) \setminus \{c_v\}.$$

Then $G >_{lex} F_k$, and since $p < c_v$, $\prod_{u \in G} u \leq n$, so $G \in \Delta([n])$. Clearly,

$$G \cap F_k = F_k \setminus \{c_v\}.$$

Now G need not be a facet, but it is contained in one, say $G \subseteq F_j$, and by (27) it follows that $F_j \geq_{lex} F_k$, whence $j < k$. Since

$$F_j \cap F_k \supseteq G \cap F_k = F_k \setminus \{c_v\},$$

and since, as noted, F_k can not be contained in another facet, we must have that

$$F_j \cap F_k = F_k \setminus \{c_v\},$$

as desired.

Case 2: Since $F_i >_{lex} F_k$ then $b_1 < c_1 < c_2 < \dots < c_s$. If all $c_1, \dots, c_s$ are prime, then $\gcd(b_1, c_v) = 1$ for $1 \leq v \leq s$. Hence

$$G = (F_k \cup \{b_1\}) \setminus \{c_1\} \in \Delta([n]).$$

As before, it suffices to note that $G >_{lex} F_k$ and that $G \cap F_k = F_k \setminus \{c_1\}$ to be able to conclude that there is some $j < k$ such that $G \subseteq F_j$,

$$F_j \cap F_k = G \cap F_k = F_k \setminus \{c_1\}.$$

□

It follows that all the homology groups of $\Delta([n])$ are torsion-free.

7. SOCLE DEGREE, THE GORENSTEIN PROPERTY, AND SYMMETRIC HILBERT FUNCTION FOR $\mathcal{A}_{[n]}$

7.1. The socle of $\mathcal{A}_{[n]}$. The following easy Lemma was proved in [18].

Lemma 7.1. *Socle($\mathcal{A}_{[n]}$) is spanned as a C -vector space by the set*

$$\begin{aligned} & \{e_k \mid 1 < k \leq n, e_k \oplus f = \mathbf{0} \text{ for all } f \text{ with } f(0) \neq 0\} = \\ & \{e_k \mid 1 < k \leq n, kp > n \text{ for all } p \in \mathbb{P} \text{ such that } \gcd(k, p) = 1\} \end{aligned} \quad (28)$$

Furthermore, the e_k 's which span the socle correspond precisely to the facets (maximal faces) $\sigma \in \Delta([n])$, $\sigma = \{p_{i_1}^{a_1}, \dots, p_{i_r}^{a_r}\}$, $k = p_{i_1}^{a_1} \cdots p_{i_r}^{a_r}$.

Theorem 7.2. *Let $\dim_{\mathbb{C}} \text{Socle}(\mathcal{A}_{[n]})$ denote the vector space dimension of the socle of $\mathcal{A}_{[n]}$. Then*

$$\lim_{n \rightarrow \infty} \frac{\dim_{\mathbb{C}} \text{Socle}(\mathcal{A}_{[n]})}{n} = 1 - \frac{1}{2} + \sum_{i=1}^{\infty} \frac{\frac{1}{p_i} - \frac{1}{p_{i+1}}}{\prod_{j=1}^i p_j} \approx 0.60771435951661818 \quad (29)$$

Proof. Let $\mathbb{N}^{++} = \{k \in \mathbb{N} \mid k > 1\}$. For all $n, k \in \mathbb{N}^+$, put

$$\begin{aligned} I_{n,0} &= \mathbb{N}^{++} \cap \left[\frac{n}{2}, n \right] \\ I_{n,k} &= \mathbb{N}^{++} \cap \left[\frac{n}{p_{k+1}}, \frac{n}{p_k} \right) \end{aligned}$$

By (28), the integers $v \in I_{n,k}$ correspond to $e_v \in \text{Socle}(\mathcal{A}_{[n]})$ whenever $p_v > n$ for all prime numbers relatively prime to v . When $2p_k > n$, $I_{n,k} = \emptyset$. For the remaining k 's, we have that $I_{n,k}$ contains approximately

$$\frac{n}{p_k} - \frac{n}{p_{k+1}}$$

integers. Of those integers, only those that are divisible by $p_1, \dots, p_k$ correspond to $e_k \in \text{Socle}(\mathcal{A}_{[n]})$. Thus, the contribution to the socle from $I_{n,k}$ is approximately

$$\frac{\frac{n}{p_i} - \frac{n}{p_{i+1}}}{\prod_{j=1}^i p_j}.$$

Clearly, all integers in $I_{n,0}$ are in the socle, which gives a contribution of approximately $\frac{n}{2}$. Furthermore, for the interval $I_{n,k}$ to contain any

integers, it must have length ≥ 1 , i.e. $\frac{n}{p_i} - \frac{n}{p_{i+1}} \geq 1$. In particular, we must have that $\frac{n}{p_i} > 1$, that is $p_i < n$. Hence, we need only consider $\pi(n)$ such intervals. For each interval $I_{n,k}$ that does contain integers, the error

$$-1 < \left(\sum_{\substack{x \in I_{n,k} \cap \mathbb{N} \\ p_1 \cdots p_k | x}} 1 \right) - \frac{\frac{n}{p_k} - \frac{n}{p_{k+1}}}{p_1 \cdots p_k} < 1.$$

Thus, we get that

$$\dim_{\mathbb{C}} \text{Socle}(\mathcal{A}_{[n]}) \approx \frac{n}{2} + \sum_{\substack{k \geq 1 \\ p_k \leq n}} \frac{\frac{n}{p_k} - \frac{n}{p_{k+1}}}{\prod_{j=1}^k p_j} \quad (30)$$

with an error $< \pi(n) \approx n/\log(n)$, from which (29) follows. $\square$

In [18] we defined the *multiplicative syzygies* of $\mathcal{A}_V$ as the kernel $K_2(V)$ of the $\mathbb{C}$ -linear map

$$\begin{aligned} \mathcal{A}_V^+ \otimes \mathcal{A}_V^+ &\rightarrow \mathcal{A}_V^+ \\ f \otimes g &\mapsto fg \end{aligned} \quad (31)$$

We call elements in $K_2(V)$ of the form $e_a \otimes e_b$ *monomial multiplicative syzygies*.

Lemma 7.3. *The monomial multiplicative syzygies of $\mathcal{A}_{[n]}^+$ correspond to the lattice points*

$$M([n]) = \{ (i, j) | 1 < i, j \leq n, ij > n \} \cup \{ (i, j) | 1 < i, j \leq n, \gcd(i, j) > 1 \} \quad (32)$$

Socle elements correspond to an integer on the x-axis such that the column supported on it is contained in $M([n])$.

Almost all syzygies are monomial, in the sense that

$$\lim_{n \rightarrow \infty} \frac{|M([n])|}{\dim_{\mathbb{C}} K_2([n])} = 1 \quad (33)$$

where $K_2([n])$ is defined as in (31).

Proof. The first two assertions are obvious. By elementary linear algebra we have that


$$\dim_{\mathbb{C}} K_2([n]) = (n-1)^2 - (n-1).$$

On the other hand,

$$|M([n])| \geq (n-1)^2 - \int_2^n \frac{ndt}{t} \geq (n-1)^2 - n \log n,$$

so (33) follows. $\square$

Below we have plotted the monomial syzygies of $\mathcal{A}_{[30]}$. One can see that 12 is in the socle.



7.2. The Gorenstein property for $\mathcal{A}_{[n]}$. A graded Artinian algebra is Gorenstein if and only if the socle is 1-dimensional. A direct computation shows that $\mathcal{A}_{[2]} \simeq \mathbb{C}[t]/(t^2)$ is Gorenstein. We note that for $n > 2$, e_{n-1} and e_n must both belong to the socle, which is then at least 2-dimensional, so then $\mathcal{A}_{[n]}$ is not Gorenstein.

7.3. Symmetric Hilbert function. A Gorenstein Artinian algebra has a symmetric Hilbert function, hence $\mathcal{A}_2(t) = 1 + t$ is symmetric. Can $\mathcal{A}_{[n]}(t)$ be symmetric for other values of n ? If $\mathcal{A}_{[n]}(t)$ symmetric, then $c_{n,\ell(n)} = c_{n,0} = 1$, which can only occur when

$$\prod_{i=1}^r p_i \leq n < p_{r+1} \prod_{i=1}^{r-1} p_i \quad (34)$$

for some r : in this case, $\ell(n) = r$ and the only integer in the interval $[1, n]$ which is the product of r primes is $\prod_{i=1}^r p_i$. Checking these intervals for $1 \leq r \leq 10$, we get the matches displayed below.

r	n	$\mathcal{A}_{[n]}(t)$
1	2	$1 + t$
2	6	$1 + 4t + t^2$
2	7	$1 + 5t + t^2$
2	8	$1 + 6t + t^2$
2	9	$1 + 7t + t^2$
3	40	$1 + 19t + 19t^2 + t^3$

Furthermore, we have [11, §§ 22.11] that the average order and the normal order of $\omega(n)$ is $\log \log n$. For (34) we have that

$$\begin{aligned}
\log \log n &< \log \log(p_{r+1} \prod_{i=1}^{r-1} p_i) \\
&< \log r \log(p_{r+1}) \\
&= \log r + \log \log p_{r+1} \\
&< \log r + \log((r+1) \log 2) \\
&= \log r + \log(r+1) + \log \log 2 \ll r/2
\end{aligned}$$

whenever r is sufficiently large. If $\mathcal{A}_{[n]}(t)$ were symmetric, it should be centred around $r/2$. Hence, for sufficiently large r , $\mathcal{A}_{[n]}(t)$ is not symmetric. There can therefore be only a finite number of n such that $\mathcal{A}_{[n]}(t)$ is symmetric. We conjecture that the examples tabulated above are in fact all such examples.

8. BASIC HOMOLOGICAL PROPERTIES

8.1. $\mathcal{A}_{[n]}$ **as a cyclic $\mathbb{C}[Y([n])]$ -module.** $\mathcal{A}_{[n]}$ is an Artinian ring and a zero-dimensional module over $\mathbb{C}[Y([n])]$, with embedding dimension $r = \pi'(n)$, and homological dimension r . Recall [11] that $r = \pi'(n) \approx \frac{n}{\log(n)}$. Furthermore, for the last Betti number we have that

$$\begin{aligned}
\beta_r(\mathbb{C}[Y([n])], \mathcal{A}_{[n]}) &= \dim_{\mathbb{C}} \text{Socle}(\mathcal{A}_{[n]}) \\
&\approx \frac{n}{2} + \sum_{\substack{k \geq 1 \\ 2p_k \leq n}} \frac{\frac{n}{p_i} - \frac{n}{p_{i+1}}}{\prod_{j=1}^i p_j} \\
&\approx 0.60771435951661818n,
\end{aligned}$$

by [19, Theorem 12.4] and Theorem 7.2. In fact, the bijection

$$\text{Tor}_r^{\mathbb{C}[Y([n])]}(\mathcal{A}_{[n]}, \mathbb{C}) \simeq \text{Socle}(\mathcal{A}_{[n]})$$

is degree-preserving, so

$$\beta_{r,j}(\mathbb{C}[Y([n])], \mathcal{A}_{[n]}) = \dim_{\mathbb{C}} \text{Socle}(\mathcal{A}_{[n]})_j.$$

For the first betti number we have

$$\beta_1(\mathbb{C}[Y([n])], \mathcal{A}_{[n]}) = \mu(A_{[n]} + B_{[n]} + C_{[n]}) = \mu(A_{[n]}) + \mu(B_{[n]}) + \mu(C_{[n]}), \quad (35)$$

the minimal number of generators of the defining ideal. Clearly

$$\mu(A_{[n]}) = r, \quad \mu(B_{[n]}) = \sum_{i=1}^r \binom{\lambda_i^{[n]}}{2} \quad (36)$$

8.2. $\mathcal{A}_{[n]}$ as a cyclic $\mathbb{C}[\overline{Y([n])}]$ -module. We can also consider $\mathcal{A}_{[n]}$ as a zero dimensional module over $\mathbb{C}[\overline{Y([n])}]$, with embedding dimension r , and infinite homological dimension. We have that

$$\beta_1(\mathbb{C}[\overline{Y([n])}], \mathcal{A}_{[n]}) = \mu(B_{[n]} + C_{[n]}) = \mu(B_{[n]}) + \mu(C_{[n]}).$$

Lemma 8.1.

$$\begin{aligned} P_{\mathbb{C}[Y([n])]}^{\mathbb{C}[\Delta([n])]}(t, \mathbf{u}) &= \text{the square-free part of } P_{\mathbb{C}[Y([n])]}^{\mathbb{C}[\overline{\Delta([n])}]}(t, \mathbf{u}) \\ &= t^{-1} \sum_{U \subset V} \mathbf{u}_U t^{|U|} \mathfrak{P}_U \\ P_{\mathbb{C}[\overline{Y([n])}]}^{\mathcal{A}_{[n]}}(t, \mathbf{u}) &= t^{-1} \sum_{U \subset V} \mathbf{v}_U t^{|U|} \mathfrak{P}_U \end{aligned} \tag{37}$$

where

$$\begin{aligned} \mathfrak{P}_U &= \sum_{i=0}^{|U|} t^{-i} \tilde{H}^i(\Delta([n])_U, \mathbb{C}) \\ \mathbf{u}_U &= \mathbf{u}_U(u_1, \dots, u_r) = \prod_{j \in U} u_j, \\ \mathbf{v}_U &= \mathbf{v}_U(u_1, \dots, u_r) = \prod_{j \in U} \frac{u_j}{1 - tu_j} \end{aligned}$$

Proof. It follows from the work of Gasharov, Peeva, and Welker [9] that

$$\beta_{i, \mathbf{a}}(\mathbb{C}[Y([n])], C[\overline{\Delta([n])}]) = \dim_{\mathbb{C}} \tilde{H}^{i-2}(1, \mathbf{y}^{\mathbf{a}})_L$$

where $(1, \mathbf{y}^{\mathbf{a}})_L$ is the order complex of the interval $(1, \mathbf{y}^{\mathbf{a}})$ in the sublattice L of Y^* generated by the minimal generators of $A_{[n]} + B_{[n]} + C_{[n]}$. Similarly,

$$\beta_{i, \mathbf{a}}(\mathbb{C}[Y([n])], C[\Delta([n])]) = \dim_{\mathbb{C}} \tilde{H}^{i-2}(1, \mathbf{y}^{\mathbf{a}})_{L'}$$

where $(1, \mathbf{y}^{\mathbf{a}})_{L'}$ is the order complex of the interval $(1, \mathbf{y}^{\mathbf{a}})$ in the sublattice L' of Y^* generated by the minimal generators of $A_{[n]} + B_{[n]} + C_{[n]}$. Since the square-free monomials form a sublattice S of Y^* , it follows that $L' = L \cap S$, hence

$$\beta_{i, \mathbf{a}}(\mathbb{C}[Y([n])], C[\Delta([n])]) = \begin{cases} 0 & \mathbf{a} \text{ not square-free} \\ \beta_{i, \mathbf{a}}(\mathbb{C}[Y([n])], C[\overline{\Delta([n])}]) & \mathbf{a} \text{ square-free} \end{cases}$$

The remaining results are immediate from the formulae in the appendix. $\square$

Theorem 8.2. *The Castelnuovo-Mumford regularity of the $\mathbb{C}[\overline{Y([n])}]$ -module $\mathcal{A}_{[n]}$ (or, equivalently, the Castelnuovo-Mumford regularity of the $\mathbb{C}[Y([n])]$ -module $\mathbb{C}[\Delta([n])]$) is $1 + v(n)$, where $v(n)$ is as defined in (14).*

Proof. It follows from (37) and (14) that the Castelnuovo-Mumford regularity is equal to

$$1 + \max \left\{ i \mid \exists U : \tilde{H}^i(\Delta([n])_U, \mathbb{C}) \neq 0 \right\} = 1 + v(n).$$

□

8.3. The Stanley-Reisner ring $\mathbb{C}[\Delta([n])]$. The Stanley-Reisner $\mathbb{C}[\Delta([n])]$ is a ring of dimension $\dim \Delta([n]) + 1 = \ell(n)$ and a $\mathbb{C}[Y([n])]$ -module of homological dimension $r - 1$. From the formulae in the appendix we get that

$$\begin{aligned} \mu(B_{[n]}) + \mu(C_{[n]}) &= \beta_1(\mathbb{C}[\overline{Y([n])}], \mathcal{A}_{[n]}) \\ &= \beta_1(\mathbb{C}[Y([n])], \mathbb{C}[\Delta([n])]) \\ &= \sum_{U \subset V} \dim_{\mathbb{C}} \tilde{H}^{|U|-2}(\Delta([n])_U, \mathbb{C}) \\ &= \sum_{i=0}^{r-2} \sum_{|U|=i+2} \dim_{\mathbb{C}} \tilde{H}^i(\Delta([n])_U, \mathbb{C}) \end{aligned} \quad (38)$$

8.4. The Koszul property. Since $A_{[n]}$ and $B_{[n]}$ are quadratic, and we know for which i the reduced simplicial homology $\tilde{H}^i(\Delta([n])_U, \mathbb{C})$ can be non-zero, we get

Corollary 8.3. *The maximal degree of a minimal generator of $C_{[n]}$, for $n > 2$, is $\max \{2, v(n)\}$. In particular, $C_{[n]}$ is quadratic for $n < 15$. Furthermore,*

$$\mu(C_{[n]}) = \sum_{i=0}^{r-2} \sum_{|U|=i+2} \dim_{\mathbb{C}} \tilde{H}^i(\Delta([n])_U, \mathbb{C}) - \sum_{i=1}^r \binom{\lambda_i^{[n]}}{2} \quad (39)$$

Proposition 8.4. *$\mathcal{A}_{[n]}$ is Koszul if and only if $n < 15$.*

Proof. $\mathcal{A}_{[n]}$ is a monomial algebra, and is quadratic iff $n < 15$. Thus by a result of Fröberg [7], $\mathcal{A}_{[n]}$ is Koszul iff $n < 15$. □

APPENDIX A. HOMOLOGICAL FORMULAE FOR STANLEY-REISNER RINGS AND INDICATOR ALGEBRAS

In this appendix, we assume that Δ is a simplicial complex on the finite set $W = \{1, \dots, r\}$. For $U \subset W$, $\Delta_U = \{\sigma \in \Delta \mid \sigma \subset U\}$; it is a simplicial complex on U . We denote by $\tilde{H}^i(\Delta_U; \mathbb{C})$ the reduced simplicial homology. Note that when $\emptyset \in \Delta$, $\tilde{H}^{-1}(\Delta_{\emptyset}; \mathbb{C}) \simeq \mathbb{C}$.

We write $S = \mathbb{C}[x_1, \dots, x_r]$, $\tilde{S} = \frac{\mathbb{C}[x_1, \dots, x_r]}{(x_1^2, \dots, x_r^2)}$, and E for the exterior algebra on the vector space of linear forms in S . Then $\mathbb{C}[\Delta]$ is a cyclic S -module, $\mathbb{C}[\overline{\Delta}]$ a cyclic $\tilde{S}$ -module, and $\mathbb{C}\{\Delta\}$ a cyclic E -module. C is a module over all these rings.

Theorem A.1 (Hochster, [12]). *Let β_i denote the i 'th Betti number of $\mathbb{C}[\Delta]$ (in a minimal free resolution of $\mathbb{C}[\Delta]$ as an S -module), and let $\beta_{i,\alpha}$ denote the corresponding multi-graded Betti number. Then*

$$\begin{aligned} \beta_{i,\alpha} &= \begin{cases} 0 & \text{if } \alpha \text{ is not square-free,} \\ \dim_{\mathbb{C}} \tilde{H}^{|U|-i-1}(\Delta_U; \mathbb{C}) & \text{if } \alpha \text{ is square-free with } \text{supp}(\alpha) = U \end{cases} \\ \beta_i &= \sum_{U \subset W} \dim_{\mathbb{C}} \tilde{H}^{|U|-i-1}(\Delta_U; \mathbb{C}) \end{aligned} \quad (40)$$

Thus the Poincaré-Betti series is

$$P_S^{\mathbb{C}[\Delta]}(t, \mathbf{u}) = \sum_{U \subset W} \prod_{j \in U} u_j \sum_{i=-\infty}^{\infty} t^i \dim_{\mathbb{C}} \left(\tilde{H}^{|U|-i-1}(\Delta_U; \mathbb{C}) \right) \quad (41)$$

Note that $\tilde{H}^j(\Delta_U; \mathbb{C}) = 0$ for $j < -1$, $j \geq n-1$, so the above sum is finite.

Theorem A.2 (Aramova-Herzog-Hibi [1]). *Let β_i denote the i 'th Betti number of $\mathbb{C}\{\Delta\}$ (in a minimal free resolution of $\mathbb{C}\{\Delta\}$ as an E -module), and let $\beta_{i,\alpha}$ denote the corresponding multi-graded Betti number. Then*

$$\beta_{i,\alpha} = \dim_{\mathbb{C}} \tilde{H}^{|\alpha|-i-1}(\Delta_U; \mathbb{C}), \quad \text{where } U = \text{supp}(\alpha) \quad (42)$$

Corollary A.3. *In the above situation,*

$$\beta_i = \sum_{U \subset W} \sum_{\ell=-1}^{|U|-1} \binom{\ell+i}{\ell+1+i-|U|} \dim_{\mathbb{C}} \tilde{H}^{|U|-i-1}(\Delta_U; \mathbb{C}) \quad (43)$$

Lemma A.4 (Sköldbberg [15]).

$$P_E^{\mathbb{C}\{\Delta\}}(t, u_1, \dots, u_r) = P_S^{\mathbb{C}[\Delta]}(t, \frac{u_i}{1-tu_1}, \dots, \frac{u_r}{1-tu_r}) \quad (44)$$

Corollary A.5.

$$P_E^{\mathbb{C}\{\Delta\}}(t, \mathbf{u}) = \sum_{U \subset W} \prod_{j \in U} \frac{u_j}{1-tu_j} \sum_{i=-\infty}^{\infty} t^i \dim_{\mathbb{C}} \left(\tilde{H}^{|U|-i-1}(\Delta_U; \mathbb{C}) \right) \quad (45)$$

Definition A.6. For $U \subset W$, we introduce the notation

$$\begin{aligned} \mathbf{u}_U &= \mathbf{u}_U(u_1, \dots, u_r) = \prod_{j \in U} u_j, \\ \mathbf{v}_U &= \mathbf{v}_U(u_1, \dots, u_r) = \prod_{j \in U} \frac{u_j}{1-tu_j}, \\ \mathfrak{P}_U &= \sum_{i=-1}^{\infty} t^{-i} \tilde{H}^i(\Delta_U, \mathbb{C}) \end{aligned} \quad (46)$$

Corollary A.7.

$$\begin{aligned}
P_S^{\mathbb{C}[\Delta]}(t, \mathbf{u}) &= t^{-1} \sum_{U \subset W} \mathbf{u}_U t^{|U|} \mathfrak{P}_U \\
P_E^{\mathbb{C}\{\Delta\}}(t, \mathbf{u}) &= t^{-1} \sum_{U \subset W} \mathbf{v}_U t^{|U|} \mathfrak{P}_U
\end{aligned} \tag{47}$$

Lemma A.8 (Sköldberg, [15]).

$$\begin{aligned}
P_{\bar{S}}^{\mathbb{C}[\bar{\Delta}]}(t, \mathbf{u}) &= P_E^{\mathbb{C}\{\Delta\}}(t, \mathbf{u}) \\
P_{\mathbb{C}[\Delta]}^{\mathbb{C}}(t, \mathbf{u}) &= P_{\mathbb{C}\{\Delta\}}^{\mathbb{C}}(t, \mathbf{u})
\end{aligned} \tag{48}$$

REFERENCES

- [1] Annetta Aramova, Jürgen Herzog, and Takayuki Hibi. Gotzman theorems for exterior algebras and combinatorics. *Journal of Algebra*, 191:174–211, 1997.
- [2] Anders Björner and Michelle L. Wachs. Shellable nonpure complexes and posets. I. *Trans. Amer. Math. Soc.*, 348(4):1299–1327, 1996.
- [3] Anders Björner and Michelle L. Wachs. Shellable nonpure complexes and posets. II. *Trans. Amer. Math. Soc.*, 349(10):3945–3975, 1997.
- [4] E. D. Cashwell and C. J. Everett. The ring of number-theoretic functions. *Pacific Journal of Mathematics*, 9:975–985, 1959.
- [5] Jean-Guillaume Dumas, Frank Heckenbach, B. David Saunders, and Volkmar Welker. Simplicial Homology. GAP package (<http://www.cis.udel.edu/~dumas/Homology/>).
- [6] S. Eliahou and M. Kervaire. Minimal resolutions of some monomial ideals. *Journal of Algebra*, 129:1–25, 1990.
- [7] Ralf Fröberg. Determination of a class of Poincaré series. *Math. Scand.*, 37(1):29–39, 1975.
- [8] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.3*, 2002. (<http://www.gap-system.org>).
- [9] Vesselin Gasharov, Irena Peeva, and Volkmar Welker. The lcm-lattice in monomial resolutions. *Math. Res. Lett.*, 6(5-6):521–532, 1999.
- [10] Daniel R. Grayson and Michael E. Stillman. Macaulay 2. Computer algebra program, available at <http://www.math.uiuc.edu/Macaulay2/>.
- [11] G. H. Hardy and E. M. Wright. *An introduction to the theory of numbers*. The Clarendon Press Oxford University Press, New York, fifth edition, 1979.
- [12] Melvin Hochster. Cohen-Macaulay rings, combinatorics, and simplicial complexes. In *Ring theory, II (Proc. Second Conf., Univ. Oklahoma, Norman, Okla., 1975)*, pages 171–223. Lecture Notes in Pure and Appl. Math., Vol. 26. Dekker, New York, 1977.
- [13] Diane MacLagan. Boolean term orders and the root system B_n . *Order*, 15(3):279–295, 1998/99.
- [14] W. Narkiewicz. On a class of arithmetical convolutions. *Colloq. Math.*, 10:81–94, 1963.
- [15] Emil Sköldberg. Monomial Golod Quotients of Exterior Algebras. *Journal of Algebra*, 218:183–189, 1998.
- [16] Jan Snellman. Truncations of the ring of number-theoretic functions. *Homology, Homotopy and Applications*, 2:17–27, 2000, math.NT/9904143.
- [17] Jan Snellman. The ring of arithmetical functions with unitary convolution: Divisorial and topological properties. *Research Reports in Mathematics*

- 4/2002, Department of Mathematics, Stockholm University, apr 2002, math.RA/0201082.
- [18] Jan Snellman. The ring of arithmetical functions with unitary convolution: General truncations. *Research Reports in Mathematics* 6/2002, Department of Mathematics, Stockholm University, may 2002, math.RA/0205242.
 - [19] Richard P. Stanley. *Combinatorics and Commutative Algebra*, volume 41 of *Progress in Mathematics*. Birkhäuser, 2 edition, 1996.
 - [20] Gérald Tenenbaum. *Introduction to analytic and probabilistic number theory*. Cambridge Univeristy Press, 1995.

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, SE-10691
STOCKHOLM, SWEDEN
E-mail address: jans@matematik.su.se